

# New England Agricultural Statistics Service

a field office of the National Agricultural Statistics Service United States Department of Agriculture

22 Bridge St, Third Floor
PO Box 1444
Concord, NH 03302

Aubrey R. Davis, Director www.usda.gov/nass

603-224-9639 Fax: 603-225-1434 nass-nh@nass.usda.gov



# Maine Potatoes Agricultural Chemical Usage, 1999

May 19, 2000

A special "THANK YOU" goes to Maine potato growers who have helped us by reporting fertilizer and pesticide application data in the Potato Production Practices Report in the Agricultural Resource Management Study. In Maine, data was collected in October and November of 1999.

## FERTILIZER AND PESTICIDES APPLIED TO FALL SEASON POTATOES IN 1999

The agricultural chemical use estimates in this report refer to on-farm use of commercial fertilizers and pesticides on targeted crops for the 1999 crop year. Farm and ranch operators were enumerated late in the growing season or after the farm operator had indicated that planned applications were completed. The data were compiled from the Agricultural Resources Management Study (ARMS), conducted primarily during the months of October-December of 1999.

Targeted crops in the 1999 ARMS include corn, upland cotton, fall potatoes, soybeans, peanuts, and sunflowers. Winter wheat was a target commodity for Indiana only.

Eleven fall potato producing States were included in the 1999 survey: The number of usable reports for each state were: Colorado (89), Idaho (342), Indiana (23), Maine (202), Michigan (101), Minnesota (63), North Dakota (49), Oregon (104), Pennsylvania (131), Washington (126), and Wisconsin (92).

Nitrogen fertilizer was applied to 100 percent of the fall potato acreage in all States, except in Colorado, Minnesota, North Dakota, and Pennsylvania. The number of nitrogen applications averaged 3.8 per acre with a total of 243.2 million pounds applied. Phosphate was applied to 98 percent of the acres in the States surveyed with a total of 190.1 million pounds being applied. Potash was applied to 88 percent of the fall potato acreage. About 163.6 million pounds of potash were applied in total.

Herbicides were applied to 93 percent of the potato acreage in 1999 in the 11 States surveyed. Coverage ranged from 100 percent in Maine, Michigan, and Oregon to 67 percent of the crop receiving herbicides in Indiana. Metribuzin was the most widely applied active ingredient and was used on 73 percent of the total acreage while EPTC was applied to 26 percent of the planted acres. Pendimethalin and Rimsulfuron were applied to 20 and 15 percent of the planted acres, respectively.

Insecticides were applied to 93 percent of the 1999 fall potato acreage. Usage ranged from treatment on 100 percent of the acres in Michigan and Wisconsin to 76

percent of the acres in Colorado. The three most common active ingredients were Imidacloprid, Methamidophos, and Phorate which were applied to 34, 29, and 23 percent of the fall potato acreage, respectively.

Fungicide treatments were applied to 95 percent of the fall potato acreage. Maine treated 100 percent of the acreage, followed closely by Michigan and North Dakota with 99 percent each. Chlorothalonil was used the most, as it was applied on 72 percent of the acreage, followed by Mancozeb on 65 percent of the fall potato acreage.

Usage of other chemicals, primarily desiccants, varied widely among the 11 States with an average of 45 percent of the fall potato acreage being treated. Diquat was the most commonly used active ingredient in the States surveyed and was applied to 38 percent of the planted area.

The use of trade names in this report is for information only and should not be construed as a recommendation by NASS.

This report is taken from the national Agricultural Chemical Usage 1999 Field Crops Summary report published by USDA's National Agricultural Statistics Service at 3 pm on May 17, 2000. The full report (about 115 pages) includes the 1999 farm use of fertilizers and pesticides applied to corn, upland cotton, fall potatoes, soybeans, peanuts, sunflowers and winter wheat

The information printed here features statistics for Maine and shows an overview of agricultural chemicals applied to fall potatoes in major potatoproducing states. The full report also shows each of the other states individually in addition to Maine.

National Reports can be ordered by calling 1-800-999-6779. How can you get these reports electronically?

- \* All national reports and state newsletters are available on the Internet.
- \* For free national e-mail reports, send a message to: usda-reports@usda.mannlib.cornell.edu and in the body, type: lists
- \* For free state newsletters, such as this, send a message to: nass-state-releases@news.usda.gov and in the body, type: subscribe new-eng-all-reports OR lists for other states.

Fall Potatoes: Fertilizer Use, Total Acreage, Percent of Acres Treated and Total Amount Applied, Eleven States Surveyed and Total, 1999

State	:	Pl anted	:	: Percent of Acres Treated and Total Applied								
State	:	Acreage	:	: Ni trogen		:	: Phosphate		: Potash			
	: 1	,000 Acres	Perd	cent 1	лі I .		Percent	Mil.	P	ercent	Mil.	
CO	:	77. 2	98	3 14	1. 6		95	13. 3		74	5. 6	
I D	:	395	100	) 9.	1.0		99	78. 5		82	42. 7	
ΙN	:	5. 2	100	) (	0. 6		100	0. 5		100	0. 5	
ME	:	65	100	) 1	1.5		100	12. 3		100	12. 4	
MI	:	48	100	) 10	). 1		98	6. 6		100	10. 0	
MN	:	70	90	9 8	3. 0		91	5. 3		91	9. 6	
ND	:	121	90	9 1!	5. 4		98	10. 9		83	9. 2	
OR	:	56	100	) 1:	3. 5		100	8. 2		91	7. 5	
PA	:	14. 5	9	7 :	2. 2		97	1.8		97	2. 0	
WA	:	170	100	5!	5. 5		99	40. 7		97	43. 7	
WI	:	86	100	) 20	). 8		100	12. 0		99	20. 4	
Total	:	1, 107. 9	100	24:	3. 2		98	190. 1		88	163. 6	

Fall Potatoes: Fertilizer Primary Nutrient Applications, Eleven States Surveyed and Total, 1999

	: Pl anted : Acreage		Appli - : cations :		: Rate ¡	per : Total Year : Applied
	: 1,000 Acres	Percent	Number	Pounds per	Acre	Mil. Lbs
Col orado:	: 77.2					
Ni trogen	:	98	6. 2	31	194	14. 6
Phosphate	:	95	1. 3	132	183	13. 3
Potash	:	74	1. 7	56	99	5. 6
I daho:	: 395					
Ni trogen	:	100	4. 5	51	231	91. 0
Phosphate	:	99	2. 3	86	200	78. 5
Potash	:	82	1. 6	79	131	42. 7
I ndi ana:	: 5.2					
Ni trogen	:	100	1. 7	68	121	0. 6
Phosphate	:	100	1. 0	103	104	0. 5
Potash	:	100	1. 0	95	97	0. 5
Mai ne:	: 65					
Ni trogen	:	100	1. 1	153	177	11. 5
Phosphate	:	100	1. 0	176	189	12. 3
Potash	:	100	1. 0	180	191	12. 4
Mi chi gan:	: 48					
Ni trogen	:	100	3. 8	55	211	10. 1
Phosphate	:	98	1. 1	120	140	6. 6
Potash	:	100	1. 3	152	210	10. 0
Mi nnesota:	: 70					
Ni trogen	:	99	2. 1	53	115	8. 0
Phosphate	:	91	1. 1	74	84	5. 3
Potash	:	91	1. 2	126	152	9. 6
North Dakota:	: 121					
Ni trogen	:	99	3. 4	37	129	15. 4
Phosphate	:	98	1. 2	75	91	10. 9
Potash	:	83	1. 2	73	92	9. 2
Oregon:	: 56					
Ni trogen	:	100	2. 4	99	242	13. 5
Phosphate	•	100	1. 6	89	147	8. 2
Potash	•	91	1. 4	100	147	7. 5
Pennsyl vani a:	: 14.5					
Ni trogen		97	1. 6	92	155	2. 2
Phosphate	•	97	1. 1	104	124	1. 8
Potash	:	97	1. 2	113	141	2. 0
Washi ngton:	: 170	,,	1. 2	110		2. 0
Ni trogen	. 170	100	3. 6	89	327	55. 5
Phosphate		99	1. 6	142	242	40. 7
Potash		97	1. 3	194	264	43. 7
Wi sconsi n:	: 86	, ,	1. 3	174	204	70.7
Ni trogen	. 55	100	4. 4	54	243	20. 8
Phosphate		100	1. 5	90	140	12. 0
Potash		99	2. 5	95	239	20. 4
Total :	: 1, 107. 9	77	۷. ن	73	237	20. 4
Ni trogen	. 1, 10/.7	100	3. 8	57	220	243. 2
•		98	3. 8 1. 7	101	220 174	243. 2 190. 1
Phosphate Potash	: :	98 88	1. 7 1. 5	101	166	163. 6
		OO	1. 5	100	100	103. 0

Fall Potatoes: Agricultural Chemical Applications, Eleven States Surveyed, 1999 1/

Agri cul tural Chemi cal	Appl i ed	: cations :	Rate per Application	: Crop Year	: Total : Applied
:	Percent	Number		per Acre	1,000 lbs
Herbi ci des:	2	1 0	0.0/	0.10	1
2, 4-D : EPTC :	2 26	1. 9 1. 0	0. 06 3. 48	0. 12 3. 69	1 1, 077
GI yphosate	3	1. 0	0. 65	0. 65	20
Li nuron :	5	1. 0	0. 63	0. 63	35
Metol achl or	16	1. 0	1. 75	1. 76	315
Metri buzi n	73	1.0	0. 42	0.45	366 170
Pendi methalin : Ri msul furon :	20 15	1. 0 1. 0	0. 80 0. 02	0. 81 0. 02	178 2
Sethoxydi m	3	1. 0	0. 29	0. 29	9
Trifluralin :	8	1. 0	0. 51	0. 51	43
Insecticides:	_				
Al di carb	5	1.0	2. 65	2. 72	141
Azinphos-methyl : Bt (Bacillus thur.)2/:	7	1. 1 1. 6	0. 42	0. 48	38
Carbaryl :	3	1. 2	0. 85	1. 02	30
Carbofuran :	10	1. 2	1. 44	1. 85	204
Cyfluthrin :	3	1. 3	0. 03	0. 04	**
Di azi non	2	1.1	1. 43	1. 71	31
Di methoate : Di sul foton :	13	1. 7 1. 1	0. 38 1. 31	0. 68 1. 49	100 19
Ethoprop :	8	1. 0	3. 83	3. 88	331
Endosul fan	16	1. 4	0. 85	1. 21	221
Esfenval erate :	16	1.4	0. 04	0.05	9
Fonofos :	1	1. 3	1. 68	2. 35	30
l mi dacl opri d Lambda-cyhal othri n	34	1. 2 1. 6	0. 12 0. 21	0. 15 0. 34	55 **
Mal athi on	*	1. 0	0. 66	0. 66	2
Methami dophos	29	1. 7	0. 91	1. 61	520
Methomyl :	*	1. 5	0. 33	0. 51	**
Methyl parathion :	1	1.7	0. 57	0. 97	12
Oxamyl Permethrin	2 8	1. 2 1. 3	0. 62 0. 13	0. 76 0. 17	13 15
Phorate :	23	1. 0	2. 74	2. 75	691
Phosmet :	4	1. 2	0. 65	0. 81	32
Piperonyl butoxide :	2	1. 9	0. 43	0. 82	22
Propargi te	4	1.1	1. 64	1. 92	76 **
Pyrethrins : Spinosad :	3	1. 6 1. 0	0. 04 0. 05	0. 06 0. 05	2
Fungi ci des:		1.0	0. 03	0.03	2
Azoxystrobin :	24	2. 5	0. 10	0. 26	69
Chl orothal oni l	72	4. 0	0. 93	3. 73	2, 960
Copper ammonium :	2	1. 9	0. 33	0.65	16
Copper hydroxide : Copper sulfate :	13	1. 6 1. 0	0. 56 1. 10	0. 94 1. 14	134 25
Cymoxani I	13	1. 7	0. 11	0. 20	29
Dimethomorph :	1	1. 3	0. 18	0. 23	3
I prodi one	7	1.0	0. 97	1. 05	87
Mancozeb :	65	3. 1 2. 5	1. 21	3.86	2, 775
Maneb : Mefenoxam :	5 20	2. 5 1. 5	1. 23 0. 11	3. 19 0. 17	190 37
Metal axyl	12	1. 4	0. 22	0. 31	40
Metiram´:	12	2. 6	1. 46	3.83	529
Propamocarb hydroch. :	1	2. 5	0. 57	1. 47	18
Sul fur :	4	1.8	2. 12	3. 89	188
Triphenyltin hydrox. : Other Chemicals:	18	2. 1	0. 14	0. 29	58
Di chl oropropene :	5	1. 0	171. 99	173. 45	10, 425
Di quat	38	1. 3	0. 34	0. 48	202
Endothall :	3	1.0	0. 79	0.86	23
Maleic hydrazide :	6	1.0	0. 97	0. 98	62
Metam-sodium :	22	1. 0 1. 0	121. 43 0. 32	121. 98 0. 32	29, 327 5
Paraquat : Sulfuric acid :	18	1.0	281. 06	293. 09	58, 893
				,_,	

<sup>\*</sup> Area applied is less than one percent.

\*\* Total applied is less than 1,000 lbs.

1/ Planted acres in 1999 for the 11 states surveyed were 1.11 million acres.

States included are CO, ID, IN, ME, MI, MN, ND, OR, PA, WA and WI.

2/ Rates and total applied are not available because amounts of active ingredient are not comparable between products.

Fall Potatoes: Pesticide Use, Total Acreage, Percent of Area Receiving Applications and Total Applied, Eleven States Surveyed and Total, 1999

: State:	: Planted	: Area Receiving and Total Applied									
3tate. :	Acreage :	Herbi ci de :		Insect	i ci de 1/:	Fungi	ci de :	Other Chemical			
:	1,000 Acres	Percent	1,000 Lbs	Percen	t 1,000 Lbs	Percen	t 1,000 Lbs	Percer	nt 1,000 Lbs		
CO : ID : IN 2/:	77. 2 395 5. 2	92	175 953 9	76 92 99	39 1, 066 2	98 92 29	387 1, 502 10	57 56	14, 056 53, 358		
ME : MI : MN :	65 48 70	100 100 86	25 101 82	97 100 91	29 52 54	100 99 93	553 609 577	24 56 16	89 137 2, 103		
ND : OR : PA :	121 56 14. 5	83 100 94	94 129 35	95 89 99	121 183 47	99 97 95	966 314 125	5 65 3	1, 315 7, 489 4		
WA : WI :	170 86	98 98	360 84	99 100	810 193	97 98	1, 206 921	75 16	19, 377 1, 104		
Total:	1, 107. 9	93	2, 047	93	2, 596	95	7, 170	45	99, 032		

<sup>1/</sup> Total Applied excludes Bt's (Bacillus thuringiensis). Total BT quantities are not available because amounts of active ingredient are not comparable between products. 2/ Insufficient reports to publish data for one or more of the pesticide classes.

Fall Potatoes: Agricultural Chemical Applications, Maine, 1999 1/

Agri cul tural : Chemi cal :	Area Appl i ed		Rate per : Application :		
:	Percent	Number	Pounds pe	er Acre	1,000 lbs
Herbi ci des:	0	1.0	0.54	0.50	
Li nuron :	8	1.0	0. 51 0. 40	0. 53 0. 41	3
Metri buzi n : Ri msul furon :	82 8	1. 0 1. 1	0. 40	0. 41	22 * *
Insecti ci des:	O	1. 1	0.02	0. 02	
Azi nphos-methyl :	3	1. 3	0. 29	0. 40	1
Bt (Bacillus thur.)2/:	*	3. 3			
Carbaryl :	1	2. 7	0. 77	2. 10	2
Di sul foton :	6	1. 4	0. 62	0. 91	2 3 2
Endosul fan :	4	1. 1	0. 59	0. 67	2
Esfenval erate :	8 90	1. 5 1. 3	0. 03 0. 09	0. 05 0. 12	7
Methami dophos :	19	1. 3	0. 61	1. 08	13
Permethrin :	ĺź	1. 5	0. 10	0. 16	1
Fungi ci des:	_				
CȟI orothal oni I :	72	7. 3	0. 62	4. 50	212
Copper hydroxide :	30	2. 1	0. 37	0. 80	16
Cymoxani I :	3	2. 7	0. 12	0. 32	1
Mancozeb : Maneb :	77 5	6. 0 6. 4	0. 97 0. 95	5. 87 6. 14	293 18
Mefenoxam :	19	1. 9	0. 95 0. 12	0. 14	3
Metalaxyl :	4	2. 1	0. 12	0. 23	1
Metiram :	2	3. 6	1. 21	4. 46	5
Triphenyltin hydrox. :	26	2. 2	0. 12	0. 26	4
Other Chemicals: :					
Di quat :	93	1. 7	0. 25	0. 43	26
Endothall :	3 16	1. 0 1. 0	0. 48 1. 29	0. 48 1. 29	14
Maleic hydrazide :	10	1. U	1. 27	1. 29	14

#### Listed below are persons within the National Agricultural Statistics Service to contact for additional information.

Jorge Garcia-Pratts, Environmental Statistician (202) 720-7492 Norman Bennett, Head, Environmental and Demographics Section (202) 720-0684 Linda Hutton, Chief, Environmental, Economics and Demographics Branch (202) 720-6146

### Listed below is the contact within the Economic Research Service for additional information.

Merritt Padgitt, Data and Survey Coordinator Resource Economic Division (202) 694-5620

AUBREY R. DAVIS, Director MARC TOSIANO, Deputy Director STATISTICIANS: Suzanne Adams

Area applied is less than one percent. Total applied is less than 1,000 lbs. Planted acres in 1999 for Maine were 65,000 acres. Rates and total applied are not available because amounts of active ingredient are not comparable between products.